



IMAGE: TICKSAFETY.COM

BROWN DOG TICK

HABITAT

They prefer to stay close to places where there is a lot of humans and animals available to feed on, such as inside homes and dog kennels.

DISEASE

Can cause Rocky Mountain Spotted Fever.



IMAGE: THE UNIVERSITY OF MAINE: TICK LAB

SQUIRREL TICK

HABITAT
They are also
frequently found
in abandoned or
seasonal buildings.

DISEASE
Can carry
Powassan virus.



IMAGE: THE UNIVERSITY OF MAINE: TICK LAB

MOUSE TICK

HABITAT
Typically only
found in the nest
of its host.

DISEASE
Can carry
Lyme Disease.



IMAGE: TICKSAFETY.COM

GULF COAST TICK

HABITAT

They are found in grass prairies and coastal uplands, like coastal banks, dunes, and sea cliffs.

DISEASE

Can cause
Spotted Fever.



IMAGE: TICKSAFETY.COM

WESTERN BLACK LEGGED TICK

HABITAT
Can be found in
vegetation in warm,
moist areas.

DISEASE
Can cause
Lyme Disease.



IMAGE: TICKSAFETY.COM

BLACK LEGGED TICK

HABITAT

can be found under leaf-litter, at the edges of wooded areas, in areas of high grass or thick vegetation, and areas where wildlife are frequent.

DISEASE

Can cause Lyme Disease.



IMAGE: TICKSAFETY.COM

GROUND HOG TICK

HABITAT

Can be found
around burrows
or dens.

DISEASE

Can cause
Powassan virus.



IMAGE: TICKSAFETY.COM

LONE STAR TICK

HABITAT

Can be found
in thick bushes
and shrubbery.

DISEASE

Can cause
monocytic
ehrlichiosis.